Content available at: <https://www.ipinnovative.com/open-access-journals>

The Journal of Community Health Management

Journal homepage: <https://www.jchm.in/>

Review Article

Effect of tobacco consumption – A review

Rakesh Kumar Pandey¹, Jeevan Josh², S C Mohapatra^{3,*}¹Dentist & MPH Scholar, Gurugram University, Gurugram, Haryana, India²Teerthanker Mahaveer Dental College, Bagadpur, Uttar Pradesh, India³SGT University, Gurugram, Haryana, India

ARTICLE INFO

Article history:

Received 28-02-2021

Accepted 01-03-2021

Available online 30-04-2021

Keywords:

Behaviour Change in Tobacco

Nicotine replacement therapy

Smoking

ABSTRACT

There are about 1.2 billion smokers worldwide of which estimated about 4 million died every year. Tobacco consumption is a common habitual concern in India; in two major Forms such as Smoking and Chewing tobaccos. Smoking includes Cigarette, Bidi Hukkah and other forms of consumption while Chewing tobaccos include Khaini, Gutka, Mawa, Dukta, Snuff, Gudakhu and other forms. The cancer reported by Chewing forms of Tobaccos are known as CIT (Chewable Indian Tobacco) Cancers. The CIT cancers more seen in India and adjacent countries like Nepal, Bangladesh, Pakistan, Sri Lanka etc, than western part of the globe. It has been calculated that 26 grams of CIT for 20 years will produce oral cancer in genetically coded people. It is a major problem in Oncology, Oral medicine as well as Public Health. Although tobacco smoking has reduced in the recent past in India, tobacco consumption / smoking is leading preventable cause of death due to cancer in males. Cigarette smoking also complicates morbidities related pulmonary conditions like Asthma, COPD, Cancer and COVID etc. A dynamic comprehensive approach is required for breaking the habit of tobacco consumption in any form. Various studies conducted in the past proposed that intensive tobacco cessation interventions involving behavioural support plays an important role in the treatment of addiction / deaddiction. Public health professionals can play an important role in creating awareness among people regarding the ill effects of tobacco through different community communication technologies. It is important that government has passed trading restrictions in order to curb the tobacco consumption through primordial prevention method. Advertisement and promotion of tobacco products are also banned throughout the country, though it continues in one form or other. Many NGO as well as Government sectors have taken initiatives at central and state level to establish tobacco cessation clinics (TCCs) in fewer instances. Behaviour Change in Tobaccoreplacement therapy, Smoking.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Tobacco comes from the leaves of *Nicotiana tabacum* which are raised, dehydrated and preventable cause of death, although it is Primarily a Risk factor. However, reduction in tobacco consumption is seen more effectively in developed countries as compared to developing countries. India is the second largest consumer and third largest producer of tobacco.¹ Annually 7,00,000 deaths are recorded in India which are mainly attributed to tobacco consumption and

is expected to rise around 1 million in next few decades.² China followed by India are the major countries which contribute to smoking risks. (Figure 1)

In India, around 35% adults use tobacco and around 25% consume smokeless tobacco, 9% only smoke cigarettes while 5% use cigarettes as well as CIT.³ In India, annual incidence of oral cancer among males is as high as 10 per 100,000. Smokers are more prone to develop respiratory, cardiovascular and peripheral vascular diseases as compared to non-smokers. Approximately, one person dies due to tobacco consumption every six seconds.^{4,5} Smoking is considered as a chronic relapsing condition,

* Corresponding author.

E-mail address: vishwamegh@gmail.com (S. C. Mohapatra).

Fig. 1: Worldwide distribution of smokers; MPOWER package, 2008

whose treatment is often difficult. As reported by American Heart Association, nicotine dependence is hardest to interrupt because its behavioural as well as pharmacological features those are equivalent to that of cocaine and heroin dependence.⁶

2. Second Hand Smoke

It is also called as passive smoking, involuntary smoking and environmental tobacco smoking. Second hand smoking is basically inhalation of tobacco smoke by persons other than the intended “active” smoker, which affect the children around (Figure 2).

Fig. 2: Diseases and adverse health effects caused by second hand smoke

3. Third Hand Smoke

Third hand smoke refers to the chemical residual of tobacco smoke built up for weeks and months that clings to clothing, cushions, furniture, wall, carpet, hair, skin, and other materials after the cigarette is extinguished. This normally affects the young ones who quire toxins by tongue while playing (Figure 3)

Fig. 3: Third hand smoking biggest risk to young children

3.1. Green Tobacco Sickness (GTS)

GTS is a form of nicotine sickness/poisoning that occurs due to nicotine dermal exposure with wet tobacco leaves. Commonly seen among people who harvest or cultivate tobacco. The overall prevalence of GTS is higher (86.2%) among tobacco cultivators. Symptoms mainly includes headache, weakness, giddiness, nausea, vomiting, abdominal pain and breathlessness.

4. Musculoskeletal effect Review of Tobacco Consumption

Tobacco consumption has many deleterious effects on the musculoskeletal system which in turn reduces the quality of life. Studies conducted by different researchers who studied the effect of smoking on musculoskeletal system has been depicted in Table 1.

Table 1: Effect of smoking on musculoskeletal system

Citations	Results
Beyth et al., 2015 ⁷	Average concentration of Bone marrow progenitor cells (BMPCs) in smokers is lower as compared to nonsmokers.
Bjarnason et al., 2009 ⁸	Bone mineral density of smokers on treatment with estradiol increases but with a less pace as compared to nonsmokers
Breitling, 2015 ⁹	The study was carried out on three different groups (i.e. never-smokers, former smokers and current smokers) and they showed an overall positive trend between calcium consumption and BMD
Callreus et al., 2013 ¹⁰	The relative fracture risk and bone mineral density is same in never-smokers and former and current smokers; however, the bone mineral density of femoral neck gets reduced as smoking increases. The bone mineral density is not significantly lower in young women of a longer duration of time or just started. In case of smokers the reduced levels of bone mineral density last only for 24 months and becomes equal to those of nonsmokers with 24 months of time.
Cangussuet al., 2012 ¹¹	There was no significant difference between the postural balance among smokers and nonsmokers, in the study. However, the risk of fall was seen to be higher among smokers than nonsmokers in case of postmenopausal women
Caram et al., 2016 ¹²	The results of the study showed that there is an inverse joint effect of smoking and COPD on health, composition of body and exertion.
Cetin et., 2009 ¹³	Smokers have higher rate of osteopenia and osteoporosis as compared to nonsmokers. Less T-scores in smokers Higher activity of antioxidant enzymes in smokers than nonsmokers
Chassanidis et al., 2012 ¹⁴	Smokers showed less reduction of expression of bone morphogenetic proteins (BMP-2, BMP-4 and BMP-6) Also non-fractured group showed higher expression of bone morphogenetic proteins than fractured group.
Dorn et al., 2013 ¹⁵	High frequency of smoking is associated with less bone mineral density of lumber spine and total hip among 13-19 age group of girls Higher symptoms of depression are associated with less BMD of lumber spine
Drage et al 2007 ¹⁶	No significant association between bone marrow density of jaws, hips etc and years of cigarette smoking exists.
Eleftheriou et al., 2013 ¹⁷	Smoking adversely affects the bone marrow density and calcaneal quantitative ultrasound High bone marrow density in case of moderate alcohol intake Physical activities improve the bone mineral density
Emaus et al., 2014 ¹⁸	Life time bone loss was higher in smokers than in nonsmokers Body mass index effects the loss of bone in case of women
Fujiyoshi et al., 2016 ¹⁹	According to this study, people who were current smokers showed lower levels of parathyroid hormone as compared to old smokers. Along with this smoker also showed lowest levels of serum calcium and excretion of calcium through urine in a day.
Kargin et al., 2016 ²⁰	Mean serum PTH level and C-terminal telopeptide levels were less in smokers than in nonsmokers.
Kassi et al., 2015 ²¹	The study revealed that smokers had comparatively less 25(OH)D level than nonsmokers.
Kaume et al., 2014 ²²	Smoking-induced bone loss was lesser in women consuming antioxidant rich berries is less than in non-consuming group.
Kleppinger et al., 2010 ²³	According to this study the weight, fat mass and functional muscle mass increased within a period of 16 months after quitting cigarette.
Lucas et al., 2012 ²⁴	Lower levels of bone mineral density was observed in girls at the age of 17 years who consumed smoking at the age of 13 and same goes with those who consumed alcohol.

Continued on next page

Table 1 continued

Myong, et al., 2013 ²⁵	A significant linear relationship exists between smoking and bone mineral density among Korean women.
Tamaki et al., 2010 ²⁶	Smokers showed less lumber spine bone mineral density as compared to smokers
Dinah et al., 2007 ²⁷	The success of operative management in case of nonunion carpal scaphoid was less in smokers as compared to nonsmokers.
Hernigou & Schuind, 2013 ²⁸	In case of both open as well as closed fracture smoking showed significant association with nonunion.
Campos et al., 2011 ²⁹	Smoking shows negative effect on expression of alveolar bone-related markers.
Adler et al., 2008 ³⁰	Smokers showed deep periodontal pockets as compared to nonsmokers.
Eren et al., 2015 ³¹	Lesser platelet-derived growth factor-AB are seen in smokers as compared to nonsmokers
Heikkinen et al., 2008 ³²	Boys and girls who smoke showed higher calculus and periodontal pocket as compared to those who do not smoke.
Hugoson & rolandsson, 2011 ³³	Smokers showed more significant association with severe periodontitis as compared to gingivitis. Snus users showed less significant association with gingivitis and periodontitis.
Torrungruang et al., 2012 ³⁴	Periodontal conditions were compromised among smokers as compared to nonsmokers.
Ata-Ali et al., 2016 ³⁵	Peri-implant parameters were compromised among smokers as compared to nonsmokers.
Rodriguez-Argueta et al., 2011 ³⁶	Strong association exists between smoking and complications in case of implants.
Romanos et al., 2013 ³⁷	As compared to smokers' long term success rate of dental implants was more in nonsmokers.
Shibli et al., 2010 ³⁸	Smokers showed less bone-to-implant contact and decreased bone density inside and outside the threaded area.
Sverzut et al., 2008 ³⁹	Smokers showed 3.2% of early implant loss in contrast to only 2.81%. nonsmokers.
Saevarsdottir et al., 2011 ⁴⁰	Current smokers showed lower response to treatment (MTX & TNF inhibitors).
Mattey et al., 2011 ⁴¹	Smokers showed high scores of Bath AS Functional Index as compared to nonsmokers.
Adedoyin et al., 2010 ⁴²	Athletes who smoke showed high rate of exertion and fatigue index as compared to nonsmoker athletes.
Kim & Kim, 2012 ⁴³	Smokers are more dependent on internal oblique and transverse abdominis muscles as compared to nonsmokers.
Kumar et al., 2010 ⁴⁴	Smokers showed reduction in lumber extension strength as compared to nonsmokers.
Rom et al., 2015 ⁴⁵	According to this study the subjects who quit smoking showed improvement in body parameters as well as strength as compared to those who continued smoking.
Blackwell et al., 2016 ⁴⁶	Smokers showed early meniscus repair failure as compared to nonsmokers.
Sanden et al., 2011 ⁴⁷	Requirement of analgesics was more in smokers The ability to walk after surgical treatment of lumber spine in smokers was more difficult to improve than nonsmokers.
Hagnas et al., 2016 ⁴⁸	Babies of smoker mothers were more associated with low aerobic fitness than nonsmoker mothers.
Holmberg et al., 2011 ⁴⁹	The bone mineral density of individuals who were exposed to second hand smoking was less than nonsmokers.
Zadzinska et al., 2016 ⁵⁰	According to this study there are more chances of having offspring with shorter leg length in parents who smoke than in parents who do not smoke.

5. Dental Health

Primarily, mouth is exposed to tobacco and most significant effects of tobacco affects the oral cavity includes precancerous lesions, oral cancers, periodontal diseases and poor wound healing. Tobacco smoke also causes tooth decay, wearing of teeth, black pigmentation of oral tissues, Gingivitis, palatal erosions, black hairy tongue, keratotic patches. Cigarette smoking during gestation has six times greater chances of having babies with birth defects like cleft lip and cleft palate formation.⁵¹ Primary caries are seen among children in case of maternal tobacco consumption.⁵²

Fig. 4: Oral ulceration due to tobacco consumption

6. Conclusion

The issue of smoking, invariably triggers oral cancer in the brain but other issues cannot be too emphatically forgotten. Carcinoma is the most obvious terrifying condition, for which tobacco is a recognized risk factor but not causative agent. Present paper has been specifically designed to draw the attention of researches to the important studies on tobacco consumption and its effect on health other than cancer, for which tobacco is a causative agent by producing biological and physiochemical changes in humans. Multiple aspects are still not touched since they were beyond the scope of the present review.

7. Source of Funding

None.

8. Conflict of Interest

The authors declare that there is no conflict of interest.

References

1. Jha P, Chaloupka FJ. Tobacco control in developing countries. Oxford University Press; 2000.
2. World Health Organization. Health topics: Tobacco. Available from: <http://www.searo.who.int/india/topics/tobacco/en/>.
3. Gajalakshmi V, Peto R, Kanaka TS, Jha P. Smoking and mortality from tuberculosis and other diseases in India: retrospective study of 43 000 adult male deaths and 35 000 controls. *Lancet*. 2003;362(9383):507–15. doi:10.1016/s0140-6736(03)14109-8.
4. Global Adult Tobacco Survey GATS India 2009-2010. Ministry of Health and [5] Family Welfare Government of India.
5. Amit S, Bhambal A, Saxena V, Sakeena B, Saxena S, Vanka A. Tobacco cessation [13] and counseling. A dentist's perspective in Bhopal city, Madhya Pradesh. *Indian J Dent Res*. 2011;22(3):400–3.
6. Marselos M, Ch F. Knowledge and attitudes of Greek pupils towards smoking. *Mat Med Greca*. 1989;17(2):101–10.
7. Beyth S, Mosheiff R, Safran O, Daskal A, Liebergall M. Surgeon General's Report 2004. Health consequences of Smoking. In: Cigarette Smoking Is Associated with a Lower Concentration of CD105(+) Bone Marrow Progenitor Cells," Bone Marrow Research; 2015. p. 1–6.
8. Bjarnason NH, Nielsen TF, Jørgensen HL, Christiansen C. The influence of smoking on bone loss and response to nasal estradiol. *Climacteric*. 2009;12:59–65. doi:10.1080/13697130802587689.
9. Breitling LP. Smoking as an Effect Modifier of the Association of Calcium Intake With Bone Mineral Density. *J Clin Endocrinol Metab*. 2015;100(2):626–35. doi:10.1210/jc.2014-2190.
10. Callreus M, McGuigan F, Akesson K. Adverse effects of smoking on peak bone mass may be attenuated by higher body mass index in young female smokers. *Calcified Tissue Int*. 2013;93(6):517–25.
11. Cangussu LM, Nahas-Neto J, Nahas EAP, Barral A, Butros DA, Uemura G. Evaluation of postural balance in postmenopausal women and its relationship with bone mineral density- a cross sectional study. *BMC Musculoskelet Disord*. 2012;13(1). doi:10.1186/1471-2474-13-2.
12. Caram L, Ferrari R, Bertani AL. Smoking and early COPD as independent predictors of body composition, exercise capacity, and health status. *PLoS ONE*. 2016;11(10):e0164290.
13. Cetin A, Muhtaroglu S, Saraymen R, Ozturk A, Muder-Ris I. Smoking-Induced Bone Defects May Be Due to Oxidative Damage in Postmenopausal Women. *Turkiye Klinikleri J Med Sci*. 2009;29(4):851–8.
14. Chassanidis CG, Malizos KN, Varitimidis S, Samara S, Koromila T, Kolia P, et al. Smoking affects mRNA expression of bone morphogenetic proteins in human periosteum. *J Bone Joint Surg*. 2012;94-B(10):1427–32. doi:10.1302/0301-620x.94b10.28405.
15. Dorn LD, Beal SJ, Kalkwarf HJ, Pabst S, Noll JG, Susman EJ. Longitudinal impact of substance use and depressive symptoms on bone accrual among girls aged 11–19 years. *J Adolesc Health*. 2013;52(4):393–9. doi:10.1016/j.jadohealth.2012.10.005.
16. Drage NA, Palmer RM, Blake G, Wilson R, Crane F, Fogelman I. A comparison of bone mineral density in the spine, hip and jaws of edentulous subjects. *Clin Oral Implant Res*. 2007;18(4):496–500. doi:10.1111/j.1600-0501.2007.01379.x.
17. Eleftheriou KI, Rawal JS, James LE, Payne JR, Loosemore M, Pennell DJ, et al. Bone structure and geometry in young men: The influence of smoking, alcohol intake and physical activity. *Bone*. 2013;52(1):17–26. doi:10.1016/j.bone.2012.09.003.
18. Emaus N, Wilsgaard T, Ahmed LA. Impacts of Body Mass Index, Physical Activity, and Smoking on Femoral Bone Loss: The Tromsø Study. *J Bone Miner Res*. 2014;29(9):2080–9. doi:10.1002/jbmr.2232.
19. Fujiyoshi A, Polgreen LE, Gross MD, Reis JP, Sidney S, Jacobs DR. Smoking habits and parathyroid hormone concentrations in young adults: The CARDIA study. *Bone Rep*. 2016;5:104–9. doi:10.1016/j.bonr.2016.04.003.
20. Kargin NC, Marakoglu K, Unlu A, Kebapcilar L, Korucu EN. Comparison of bone turnover markers between male smoker and non-smoker. *Acta Med Mediterr*. 2016;32(2):317–23.

21. Kassi EN, Stavropoulos S, Kokkoris P, Galanos A, Moutsatsou P, Dimas C, et al. Smoking is a significant determinant of low serum vitamin D in young and middle-aged healthy males. *Hormones (Athens)*. 2015;14(2):241–50.
22. Kaume L, Gbur EE, DiBrezza R, Howard LR, Devareddy L. Antioxidant-rich berries exert modest bone protective effects in postmenopausal smokers without improving biomarkers of bone metabolism. *J Funct Foods*. 2014;9(1):202–10. doi:10.1016/j.jff.2014.04.025.
23. Kleppinger A, Litt MD, Kenny AM, Oncken CA. Effects of Smoking Cessation on Body Composition in Postmenopausal Women. *J Women's Health*. 2010;19(9):1651–7. doi:10.1089/jwh.2009.1853.
24. Lucas R, Fraga S, Ramos E, Barros H. Early Initiation of Smoking and Alcohol Drinking as a Predictor of Lower Forearm Bone Mineral Density in Late Adolescence: A Cohort Study in Girls. *PLoS ONE*. 2012;7(10). doi:10.1371/journal.pone.0046940.
25. Myong JP, Kim HR, Choi SE, Koo JW. Dose-related effect of urinary cotinine levels on bone mineral density among Korean females. *Osteoporos Int*. 2013;24(4):1339–46. doi:10.1007/s00198-012-2107-6.
26. Tamaki J, Iki M, Sato Y. Smoking among premenopausal women is associated with increased risk of low bone status: The JPOS Study. *J Bone Miner Metab*. 2010;28(3):320–7.
27. Dinah AF, Vickers RH. Smoking increases failure rate of operation for established non-union of the scaphoid bone. *Int Orthop*. 2007;31(4):503–5. doi:10.1007/s00264-006-0231-7.
28. Hernigou J, Schuind F. Smoking as a predictor of negative outcome in diaphyseal fracture healing. *Int Orthop*. 2013;37(5):883–7. doi:10.1007/s00264-013-1809-5.
29. Campos JMD, Prati A, Cirano FR, Pimentel S, Pastore GP, Pecorari VG, et al. Smoking Modulates Gene Expression of Type I Collagen, Bone Sialoprotein, and Osteocalcin in Human Alveolar Bone. *J Oral Maxillofac Surg*. 2015;73(11):2123–31.
30. Adler L, Modin C, Friskopp J, Jansson L. Relationship between smoking and periodontal probing pocket depth pro-file. *Swedish Dent J*. 2008;32(4):157–63.
31. Eren G, Turkolü HO, Atmaca H, Atilla FG. Evaluation of GCF MMP-1, MMP-8, TGF- β 1, PDGF-AB, and VEGF levels in periodontally healthy smokers. *Turk J Med Sci*. 2015;45(4):850–6. doi:10.3906/sag-1405-112.
32. Heikkinen AM, Pajukanta R, Pitkaniemi J, Broms U, Sorsa T, Koskenvuo M, et al. The Effect of Smoking on Periodontal Health of 15- to 16-Year-Old Adolescents. *J Periodontol*. 2008;79(11):2042–7. doi:10.1902/jop.2008.080205.
33. Hugoson A, Rolandsson M. Periodontal disease in relation to smoking and the use of Swedish snus: Epidemiological studies covering 20 years. *J Clin Periodontol*. 1983;38(9):809–16.
34. Torrungruang K, Gongsakdi V, Laohaviraphab L, Likittanasombat K, Ratanachaiwong W. Association between cigarette smoking and the intraoral distribution of periodontal disease in Thai men over 50 years of age. *J Investig Clin Dent*. 2012;3(2):135–41. doi:10.1111/j.2041-1626.2011.00105.x.
35. Ata-Ali J, Flichy-Fernández AJ, Alegre-Domingo T, Ata-Ali F, Peñarocha-Diogo M. Impact of heavy smoking on the clinical, microbiological and immunological parameters of patients with dental implants: a prospective cross-sectional study. *J Investig Clin Dent*. 2016;7(4):401–9. doi:10.1111/jicd.12176.
36. Rodriguez-Argueta OF, Figueiredo R, Valmaseda-Castellon E, Gay-Escoda C. Postoperative Complications in Smoking Patients Treated With Implants: A Retrospective Study. *J Oral Maxillofac Surg*. 2011;69(8):2152–7. doi:10.1016/j.joms.2011.02.082.
37. Romanos GE, Gaertner K, Aydin E, Nentwig GH. Long-Term Results After Immediate Loading of Platform-Switched Implants in Smokers Versus Nonsmokers with Full-Arch Restorations. *Int J Oral Maxillofac Implants*. 2013;28(3):841–5. doi:10.11607/jomi.3223.
38. Shibli JA, Piattelli A, Iezzi G. Effect of smoking on early bone healing around oxidized surfaces: a prospective, controlled study in human jaws. *J Periodontol*. 2010;81(4):575–83.
39. Sverzut AT, Stabile GAV, Moraes M, Mazzonetto R, Moreira RWF. The Influence of Tobacco on Early Dental Implant Failure. *J Oral Maxillofac Surg*. 2008;66(5):1004–9. doi:10.1016/j.joms.2008.01.032.
40. Saevardottir S, Wedren S, Seddighzadeh M. Patients with early rheumatoid arthritis who smoke are less likely to respond to treatment with methotrexate and tumor necrosis factor inhibitors: observations from the epidemiological investigation of rheumatoid arthritis and the Swedish rheumatology register cohorts. *Arthritis Rheumatol*. 2011;63(1):26–36.
41. Matthey DL, Dawson SR, Healey EL, Packham JC. Relationship Between Smoking and Patient-reported Measures of Disease Outcome in Ankylosing Spondylitis. *J Rheumatol*. 2011;38(12):2608–15. doi:10.3899/jrheum.110641.
42. Adedoyin RA, Mbada CE, Odiachi AM, Adegoke BOA, Awotidebe TO. Differences in back extensor muscles fatigability for smoking and non-smoking athletes. *Isokinetics Exerc Sci*. 2010;18(3):149–55. doi:10.3233/ies-2010-0377.
43. Kim LJ, Kim N. Difference in lateral abdominal muscle thickness during forceful exhalation in healthy smokers and non-smokers. *J Back Musculoskeletal Rehabil*. 2012;25(4):239–44.
44. Kumar GP. Assessment of fatigue susceptibility of the lumbar extensor muscles amongst smokers. *Biomed Eng*. 2010;22(4):321–6.
45. Rom O, Reznick AZ, Keidar Z, Karkabi K, Aizenbud D. Smoking cessation-related weight gain-beneficial effects on muscle mass, strength and bone health. *Addiction*. 2015;110(2):326–35. doi:10.1111/add.12761.
46. Blackwell R, Schmitt LC, Flanagan DC, Magnussen RA. Smoking increases the risk of early meniscus repair failure. *Knee Surg Sports Traumatol Arthrosc*. 2016;24(5):1540–3. doi:10.1007/s00167-016-4002-8.
47. Sanden B, Forst P, Michaelsson K. Smokers show less improvement than nonsmokers two years after surgery for lumbar spinal stenosis: a study of 4555 patients from the Swedish spine register. *Spine J*. 2011;36(13):1059–64.
48. Hagnäs MP, Cederberg H, Jokelainen J, Mikkola I, Rajala U, Keinänen-Kiukkaanniemi S. Association of maternal smoking during pregnancy with aerobic fitness of offspring in young adulthood: a prospective cohort study. *Int J Obstet Gynaecol*. 2016;123(11):1789–95. doi:10.1111/1471-0528.13789.
49. Holmberg T, Bech M, Curtis T, Juel K, Grønbaek M, Brixen K. Association between passive smoking in adulthood and phalangeal bone mineral density: results from the KRAM study—the Danish Health Examination Survey 2007–2008. *Osteoporos Int*. 2007;22(12):2989–99.
50. Zdzienicka E, Kozielec S, Borowska-Strugińska B, Rosset I, Sitek A, Lorkiewicz W. Parental smoking during pregnancy shortens offspring's legs. *HOMO*. 2016;67(6):498–507. doi:10.1016/j.jchb.2016.06.002.
51. Hiremath SS. Textbook of preventive and community dentistry. Elsevier; 2011.
52. Gordon JS, Andrews JA, Crews KM, Payne TJ, Severson HH. The 5A's vs 3A's plus proactive quitline referral in private practice dental offices: preliminary results. *Tobacco Control*. 2007;16(4):285–8. doi:10.1136/tc.2007.020271.

Author biography

Rakesh Kumar Pandey, MPH Scholar

Jeevan Josh, MDS Scholar

S C Mohapatra, Former HOD Community Medicine, IMS, BHU; Dean Medical College; Dean Academic affairs; Presently Academic Adviser
<https://orcid.org/0000-0002-9605-0867>

Cite this article: Pandey RK, Josh J, Mohapatra SC. Effect of tobacco consumption – A review. *J Community Health Manag* 2021;8(1):6–11.